

Choptank

Animal Waste Management Systems: Livestock

TS Goal: 19 systems

Definition: Systems for the proper handling, storage, and use of waste generated by confined animal facilities. These include ponds, lagoons, and tanks for liquid waste, and sheds or pits for solid waste.

Applied to: manure acres (stored manure)

Nitrogen Efficiency: 75% reduction

Phosphorus Efficiency: 75% reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - High

Phosphorus - High




1998 Progress for Animal Waste Management Systems: Livestock
(as a percentage of TS goal, labeled units are systems)

Implementation Record (systems)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Animal Waste Management Systems: Poultry

TS Goal: 30 systems

Definition: Systems for the proper handling, storage, and use of waste generated by confined animal facilities. These include ponds, lagoons, and tanks for liquid waste, and sheds or pits for solid waste.

Applied to: manure acres (stored manure)

Nitrogen Efficiency: 14% reduction

Phosphorus Efficiency: 14% reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Animal Waste Management Systems: Poultry
(as a percentage of TS goal, labeled units are systems)

Implementation Record (systems)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Conservation Tillage

TS Goal: 27,134 acres

Definition: A process that uses tillage equipment to seed the crop directly into the vegetative cover or crop residue on the surface, with minimal soil disturbance.

Applied to: hitill and lotill land

Nitrogen Efficiency: land conversion reduction

Phosphorus Efficiency: land conversion reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - High

Phosphorus - High




1998 Progress for Conservation Tillage
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Cover Crops

TS Goal: 50,586 acres

Definition: Small grains planted in September or early October on land otherwise fallow with no fertilizer applied. This practice reduces nitrate leaching losses during the winter, and also reduces erosion.

Applied to: hitill and lotill land

Nitrogen Efficiency: 59% reduction

Phosphorus Efficiency: 44% reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:

Nitrogen - High


Phosphorus - High




1998 Progress for Cover Crops
(as a percentage of TS goal, labeled units are acres)

Implementation Record

(acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Nutrient Management Plan Implementation

TS Goal: 150,249 acres

Definition: A comprehensive plan to manage the amount, placement, timing and application of animal waste, fertilizer, sludge, or other plant nutrients.

Applied to: cropland

Nitrogen Efficiency: model-derived reduction

Phosphorus Efficiency: model-derived reduction


BMP Rankings:

Ability to Reduce Nutrients: Medium

Importance in Statewide Strategy:


Nitrogen - High

Phosphorus - High




1998 Progress for Nutrient Management Plan Implementation
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Retirement of Highly Erodible Land

TS Goal: 474 acres

Definition: An accelerated application of practices used in SCWQPs on lands with a high potential for soil loss.

Applied to: cropland and pasture

Nitrogen Efficiency: land conversion reduction

Phosphorus Efficiency: land conversion reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Retirement of Highly Erodible Land
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Runoff Control

TS Goal: 2 systems

Definition: Systems for the proper handling, storage, and use of waste generated by confined animal facilities. These include ponds, lagoons, and tanks for liquid waste, and sheds or pits for solid waste.

Applied to: manure

Nitrogen Efficiency: 10% reduction

Phosphorus Efficiency: 10% reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Runoff Control
(as a percentage of TS goal, labeled units are systems)

Implementation Record (systems)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

SCWQP Implementation and Treatment of Highly Erodible Land

TS Goal: 36,611 acres

Definition: A comprehensive plan addressing natural resource management of farmland directed toward the control of erosion and sediment loss, and management of animal waste or agricultural chemicals.

Applied to: cropland and pasture

Nitrogen Efficiency: 11% reduction

Phosphorus Efficiency: 21% reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:


Nitrogen - High

Phosphorus - High




1998 Progress for SCWQP Implementation and Treatment of Highly Erodible Land
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Forest Conservation

TS Goal: 167 acres

Definition: Implementation of the Forest Conservation Act, which requires the retention of a portion of forested lands on any newly developed site.

Applied to: urban land

Nitrogen Efficiency: land conversion reduction

Phosphorus Efficiency: land conversion reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Forest Conservation
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Forest Harvesting Practices

TS Goal: 1,200 acres

Definition: Application of regulatory and voluntary best management practices applied to timber harvests, including erosion and sediment control and streamside management zones.

Applied to: forest land

Nitrogen Efficiency: 50% reduction

Phosphorus Efficiency: 50% reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:

Nitrogen - Low

Phosphorus - Low



1998 Progress for Forest Harvesting Practices
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)

Annual

Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Forested Buffers

TS Goal: 132 acres

Definition: A linear strip of forest along rivers and streams that filters nutrients and sediment and enhances stream habitat.

Applied to: cropland and pasture

Nitrogen Efficiency: 56% reduction

Phosphorus Efficiency: 70% reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Forested Buffers
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Grassed Buffers

TS Goal: 812 acres

Definition: A linear strip of grass along rivers and streams that filters nutrients and sediment and enhances stream habitat.

Applied to: cropland and pasture

Nitrogen Efficiency: 41% reduction

Phosphorus Efficiency: 53% reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Grassed Buffers
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Marine Pumpouts (installation)

TS Goal: 8 marinas

Definition: A facility sited at marinas for pumping sewage from boat holding tanks to a dockside storage facility.

Applied to:

Nitrogen Efficiency: pound reduction

Phosphorus Efficiency: pound reduction


BMP Rankings:

Ability to Reduce Nutrients: High

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Marine Pumpouts (installation)
(as a percentage of TS goal, labeled units are marinas)

Implementation Record (marinas)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Nonstructural Shore Erosion Control

TS Goal: 6,560 linear feet

Definition: A practice for stabilizing eroding shorelines by establishing marsh grasses; suitable for sites with lower wave energy. Also creates wetland habitat.

Applied to:

Nitrogen Efficiency: 66 lbs per l.f. reduction

Phosphorus Efficiency: 40 lbs per l.f. reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Nonstructural Shore Erosion Control
(as a percentage of TS goal, labeled units are linear feet)

Implementation Record (linear feet)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Structural Shore Erosion Control

TS Goal: 5,460 linear feet

Definition: A practice for stabilizing eroding shorelines using stone riprap or timber bulkheads. Suitable for sites with high wave energy.

Applied to:

Nitrogen Efficiency: 101 lbs per l.f. reduction

Phosphorus Efficiency: 67 lbs per l.f. reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Medium




1998 Progress for Structural Shore Erosion Control
(as a percentage of TS goal, labeled units are linear feet)

Implementation Record (linear feet)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Tree Planting

TS Goal: 990 acres

Definition: Includes any tree plantings on any site except those along rivers and streams.

Applied to: urban land

Nitrogen Efficiency: 0 reduction

Phosphorus Efficiency: 0 reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:


Nitrogen - Low

Phosphorus - Low




1998 Progress for Tree Planting
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Enhanced Stormwater Management

TS Goal: 2,536 acres

Definition: The regulatory requirement for the control of Stormwater on all new development, including maintenance on new and existing facilities. Enhancements emphasize water quality controls in addition to water quantity controls.

Applied to: urban land

Nitrogen Efficiency: 33% reduction

Phosphorus Efficiency: 46% reduction


BMP Rankings:

Ability to Reduce Nutrients: Medium

Importance in Statewide Strategy:


Nitrogen - High

Phosphorus - Medium




1998 Progress for Enhanced Stormwater Management
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Erosion and Sediment Control

TS Goal: 362 acres

Definition: The regulatory requirement for erosion and sediment control on all new development over 5,000 square feet. Reduces the high nutrient and suspended sediment loads during the transitory construction phase.

Applied to: urban land

Nitrogen Efficiency: 33% reduction

Phosphorus Efficiency: 50% reduction


BMP Rankings:

Ability to Reduce Nutrients: Medium

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - Low




1998 Progress for Erosion and Sediment Control
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Septic Connections

TS Goal: 125 systems

Definition: The connection of failing septic systems to sewer lines.

Applied to: urban land

Nitrogen Efficiency: 55% reduction

Phosphorus Efficiency: no reduction


BMP Rankings:

Ability to Reduce Nutrients: Medium

Importance in Statewide Strategy:


Nitrogen - Medium

Phosphorus - n/a




1998 Progress for Septic Connections
(as a percentage of TS goal, labeled units are systems)

Implementation Record (systems)



Annual



Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Septic Pumping

TS Goal: 60 systems

Definition: Pumping individual septic systems once every three years, the average routine maintenance of these systems.

Applied to: urban land

Nitrogen Efficiency: 5% reduction

Phosphorus Efficiency: no reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:

Nitrogen - Low

Phosphorus - n/a



1998 Progress for Septic Pumping
(as a percentage of TS goal, labeled units are systems)

Implementation Record (systems)

Annual

Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Stormwater Management Conversion

TS Goal: 19 acres

Definition: Conversion of dry ponds for Stormwater management to extended detention or retention facilities which are more effective at nutrient removal.

Applied to: urban land

Nitrogen Efficiency: 33% reduction

Phosphorus Efficiency: 46% reduction


BMP Rankings:

Ability to Reduce Nutrients: Medium

Importance in Statewide Strategy:

Nitrogen - Low

Phosphorus - Low



1998 Progress for Stormwater Management Conversion
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)

Annual

Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Stormwater Management Retrofits

TS Goal: 195 acres

Definition: Construction of Stormwater facilities on lands previously developed without such facilities.

Applied to: urban land

Nitrogen Efficiency: 33% reduction

Phosphorus Efficiency: 46% reduction


BMP Rankings:

Ability to Reduce Nutrients: Medium

Importance in Statewide Strategy:

Nitrogen - Low

Phosphorus - Low



1998 Progress for Stormwater Management Retrofits
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)

Annual

Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.

Choptank

Urban Nutrient Management

TS Goal: 498 acres

Definition: A public education program to reduce excess lawn fertilizer use, targeted at suburban residents and businesses.

Applied to: urban land

Nitrogen Efficiency: 17% reduction

Phosphorus Efficiency: 22% reduction


BMP Rankings:

Ability to Reduce Nutrients: Low

Importance in Statewide Strategy:

Nitrogen - Medium

Phosphorus - Low



1998 Progress for Urban Nutrient Management
(as a percentage of TS goal, labeled units are acres)

Implementation Record (acres)

Annual

Cumulative

NOTES: (1) Implementation data is organized by State fiscal year (i.e. 1994 = July 1, 1993 - June 30, 1994). (2) Most implementation data is based on State program activities, funding and/or requirements. (3) BMP definitions are from "Maryland's Tributary Strategies for Nutrient Reduction" report, 1995. (4) BMP implementation data is from the Maryland Interagency Tracking Subcommittee. (5) Efficiencies are average statewide efficiencies. (6) Ongoing improvements to the Watershed Model shift the expected reductions of BMPs with efficiencies listed as "land use conversions" slightly.